

HELAGO-CZ, s.r.o. Commercial Register maintained by the Regional Court in Hradec Králové Section C, File 17879 Kladská 1082 500 03 Hradec Králové 3 Company ID: 25 96 39 61, VAT: CZ 25963961

Phone: 495 220 229, 495 220 394 Fax: 495 220 154 GSM gate: 602 123 096 E-mail: info@helago-cz.cz Web: http://www.helago-cz.cz 61140 - Midline Perineal Repair Trainer

Order code: 4104.61140



Cena bez DPH 896,00 Eur Price with VAT 1.084,16 Eur

Parameters

Obstetrics, gynecology - filter Common gynecological and obstetric actions

Quantitative unit ks

OVERVIEW

- Anatomically accurate model for teaching perineal suturing
- Model represents a 2nd degree midline vaginal tear
- Illustration pack representing each degree of perineal trauma

REALISM

- All essential layers of tissue can accept repeated sutures
- Soft skin with a similar drag and strength to human skin

VERSATILITY

- · Can be used alongside PROMPT Flex Birthing Simulator, by placing the trainer in front, to add further realism
- Quick and easy to set up during training sessions
- Clamps are provided to secure the Midline Perineal Repair Trainer to the bench top
- Trainer is compact and easily portable

CLEANING

- Should trainer require cleaning, use a soft damp cloth on the plastic parts
- Do not get material section of this trainer wet

SAFETY

- This product is latex free
- Do not perform procedures without the trainer securely clamped to a work bench or training station
- Do not introduce liquid or gel to the trainer
- Ensure care when removing sutures using small sharp scissors or a scalpel and take care to only sever practice sutures and not integral stitching

SIMULATED PATIENT

- Perineal Repair Trainers can be used with a simulated patient
- Used in conjunction with PROMPT Flex birthing models with a simulated patient

ANATOMY

- Vaginal mucosa
- Hymenal remnants
- Posterior fourchette
- Perineal muscles (bulbocavernosus and transverse presented together)
- Perineal epidermis and dermis
- Anal sphincter and mucosa

SKILLS GAINED

- · Recognition of internal anatomy
- Tissue and needle handling
- Perform mock infiltration (administering local anaesthesia, we recommend this is done "dry", without fluid)
- Deep musculature, subcuticular and superficial suturing
- Repair of 2nd degree perineal tear
- Pill rolling technique to determine classification of injury