



HELAGO-CZ, s.r.o.  
Commercial Register maintained by the Regional Court in Hradec  
Králové  
Section C, File 17879  
Kladská 1082  
500 03 Hradec Králové 3  
Company ID: 25 96 39 61, VAT: CZ 25963961  
Phone: 495 220 229, 495 220 394  
Fax: 495 220 154  
GSM gate: 602 123 096  
E-mail: [info@helago-cz.cz](mailto:info@helago-cz.cz)  
Web: <http://www.helago-cz.cz>

---

**MP1810 - Lower Limb Musculature**  
Order code: **4003.MP1810**



Cena bez DPH

2.350,00 Eur

Price with VAT

2.843,50 Eur

Parameters

Quantitative unit

ks

This 3D printed specimen preserves a superficial dissection of the lower limb musculature from the mid-thigh to mid-leg, as well as nerves and vessels of the popliteal fossa. The insertions of the muscles of the anterior, middle and posterior compartments of the thigh are visible, including the pes anserinus medially and the iliotibial tract laterally. The capsule of the knee joint has been opened anterior to demonstrate the menisci and the tibial and fibular collateral ligaments. The superficial muscles of the leg are visible, with the anterior and lateral compartment muscles highlighted deep to the crural fascia. Within the popliteal fossa, both the popliteal artery and vein are visible descending from the adductor hiatus. The sciatic nerve is seen bifurcating in the fossa, with the common fibular, tibial, and sural nerves preserved.

The proximal cross-section provides a view of the distal thigh musculature of the anterior, medial and posterior compartments. The femoral artery and vein and saphenous nerve are visible within the adductor canal. The sciatic nerve and perforating branches of the profunda femoris artery (and accompanying veins) are visible in the posterior compartment. In the distal cross-section, muscles of the anterior, lateral and posterior compartment are visible. The superficial and deep fibular nerves are visible in the anterior and lateral compartments, respectively. In the anterior compartment, the deep fibular nerve is adjacent to the anterior tibial artery and veins. In the posterior compartment, deep to soleus, both the posterior tibial and fibular arteries and veins are highlighted near the tibial nerve.